

CONTENTS

Regular papers

- Studies on molecular weights of polysaccharides of *Auricularia auricula-judae*
L. Zhang, L. Yang, Q. Ding and X. Chen (Wuhan, China) 1

- Synthesis and conformational analysis of the plant hormone (auxin) related 2-(indol-3-yl)ethyl and 2-phenylethyl β -D-xylopyranosides and their 2,3,4-tri-O-acetyl derivatives
S. Tomić (Zagreb, Croatia), B.P. van Eijck (Utrecht, The Netherlands), B. Kojić-Prodić (Zagreb, Croatia), J. Kroon (Utrecht, The Netherlands), V. Magnus, B. Nigović, G. Laćan, N. Ilić (Zagreb, Croatia), H. Duddeck (Hannover, Germany) and M. Hiegemann (Bochum, Germany) 11

- Structure determination of galacto-oligosaccharides by pyridylamination and NMR spectroscopy
K. Kimura, K. Matsumoto, C. Ishihara, K. Harada and A. Miyagi (Tokyo, Japan) 33

- Synthesis, characterization, and reverse-micellar studies of some *N*-substituted derivatives of 6-amino-6-deoxy-1,2-O-isopropylidene-D-glucose
L. Sharma and S. Singh (Amritsar, India) 43

- Synthesis of a thio analogue of *n*-propyl kojibioside, a potential glucosidase inhibitor
J.S. Andrews and B.M. Pinto (Burnaby, Canada) 51

Notes

- The Resolution of oligosaccharides by high pH anion exchange chromatography
J.M. McGuire, M.A. Elliott, H.G. Elliott and K.D. Smith (Glasgow, UK) 63

- A new synthesis of sucrose 6'-phosphate
K.B. Kim and E.J. Behrman (Columbus, OH, USA) 71

- Synthesis of (3'*R*)-3'-deoxy-3'-C-nitromethylthymidine
Y. Takahashi, Y. Honda and T. Tsuchiya (Kawasaki, Japan) 77

- Synthesis of a partially protected 1D-6-O-(2-azido-2-deoxy- α -D-glucopyranosyl)-myo-inositol: a useful precursor of glycosylphosphatidylinositols and related compounds
S. Cottaz, J.S. Brimacombe and M.A.J. Ferguson (Dundee, UK) 85

- A novel synthesis of L-fucose from D-galactose
S. Sarbajna, S.K. Das and N. Roy (Calcutta, India) 93

- Identification of *N*-acetylglucosamine and 4-O-[1-carboxyethyl]mannose in the exopolysaccharide from *Cyanospora capsulata*
D. Garozzo, G. Impallomeni, E. Spina, L. Sturiale (Catania, Italy), A. Cesàro and P. Cescutti (Trieste, Italy) 97

Gel-to-sol transition of ^{13}C -labeled (1 \rightarrow 3)- β -D-glucan, ^{13}C -SSG, assessed by ^{13}C nuclear magnetic resonance (^{13}C -NMR) spectroscopy M. Suda, N. Ohno, Y. Adachi and T. Yadomae (Tokyo, Japan)	107
<i>Preliminary communication</i>	
Binding of Maillard products to proteins: formation of pyrrole carbimines J. Nissl, M. Pischetsrieder (München, Germany), E. Klein (Stuttgart, Germany) and T. Severin (München, Germany).	C1
<i>Announcement</i>	C7
<i>Regular papers</i>	
Circular dichroism of the O-specific polysaccharide of <i>Vibrio cholerae</i> O1 and some related derivatives S. Bystrický, S.C. Szu, M. Gotoh and P. Kováč (Bethesda, MD, USA)	115
Enzymatic synthesis of mannosides and mannitrioses by reverse hydrolysis using α -mannosidase from <i>Aspergillus niger</i> K. Ajsaka, I. Matsuo, M. Isomura, H. Fujimoto, M. Shirakabe and M. Okawa (Odawara, Japan) . .	123
Utility of non-covalent complexes in the matrix-assisted laser desorption ionization mass spectrometry of heparin-derived oligosaccharides P. Juhasz and K. Biemann (Cambridge, MA, USA).	131
Bromination of regenerated chitin with <i>N</i> -bromosuccinimide and triphenylphosphine under homogeneous conditions in lithium bromide- <i>N,N</i> -dimethylacetamide H. Tseng, K. Furuhata and M. Sakamoto (Tokyo, Japan).	149
Preparation of α - and β -dienyl glycosides used as dienes in aqueous Diels-Alder reactions. Influence of the carbohydrate moiety on the thermodynamics of the reaction A. Lubineau, H. Bienaymé and Y. Queneau (Orsay, France).	163
Synthesis of photolabile mono- and di-valent α -D-mannoside-6-phosphates as chemically modifying probes for mannose-6-phosphate-receptors J. Lehmann, F. Schweizer and U.P. Weitzel (Freiburg, Germany)	181
Structure of an acidic microcapsular glycan from the reference strain (C.D.C. 866-57) for <i>Serratia marcescens</i> serogroup O1 A. Johansson, G. Widmalm (Stockholm, Sweden), P.-E. Jansson (Huddinge, Sweden) and S.G. Wilkinson (Hull, UK)	191
<i>Notes</i>	
Synthesis of 4-epi-2-deoxy-2- H_{eq} - <i>N</i> -acetylneuraminic acid and 2,4-dideoxy-2- H_{eq} - <i>N</i> -acetylneuraminic acid B.P. Bandgar (Maharashtra, India) and E. Zbiral (Vienna, Austria)	201
Synthesis of substituted 2,6-dioxabicyclo[3.1.1]heptanes: 1,3-anhydro-2,4-di- <i>O</i> -benzyl-6-deoxy- β -L-talopyranose Z. Gan and F. Kong (Beijing, China)	211
Crystal-solution structural equivalence of methyl 3,6-anhydroxyranosides S.E. Schafer, E.S. Stevens (Binghamton, NY, USA) and M.K. Dowd (New Orleans, LA, USA) . . .	217

Characterisation of the endo-polygalacturonase-resistant region of the pectin from <i>Bupleurum falcatum</i> L.—a polysaccharide with an active function in clearance of immune complexes T. Matsumoto, M. Hirano, H. Kiyohara and H. Yamada (Tokyo, Japan).	221
<i>Preliminary communications</i>	
Total synthesis of sulfated Le ^a pentaosyl ceramide A. Endo, M. Iida, S. Fujita, M. Numata, M. Sugimoto and S. Nunomura (Saitama).	C9
A total synthesis of glycononaosyl ceramide with a sialyl dimeric Le ^x sequence M. Iida, A. Endo, S. Fujita, M. Numata, Y. Matsuzaki, M. Sugimoto, S. Nunomura and T. Ogawa (Saitama, Japan)	C15
<i>Book reviews</i>	C21
<i>Author index</i>	C25
<i>Keyword index</i>	C27
<i>Contents</i>	C31